



Osmotic coefficients

Nielsen, Ole Faurskov; Bilde, Merete; Kjærgaard, Henrik Grum

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record


Citation for published version (APA):
Nielsen, O. F., Bilde, M., & Kjærgaard, H. G. (2013). *Osmotic coefficients*. Abstract from European Conference on the Spectroscopy of Biological Molecules 15, Oxford, United Kingdom.

European Conference on the Spectroscopy of Biological Molecules 15

25th – 30th August 2013
St Catherine's College
The University of Oxford
Oxford, UK

ECSBM 15

where Spectroscopy and Biology meet



European Conference on the Spectroscopy of Biological Molecules

Oxford 25-30 August 2013



Science & Technology
Facilities Council

OSMOTIC COEFFICIENTS

O. F. Nielsen¹, M. Bilde², H.G. Kjaergaard¹¹Department of Chemistry, University of Copenhagen, Universitetsparken 5, Copenhagen, Denmark²Department of Chemistry, University of Aarhus, Langelandsgade 140, Aarhus, Denmark
ofn@chem.ku.dk

To preserve the overall shape of living cells the osmolarity/osmolality plays an important role.¹ The osmolarity of the extracellular fluid is usually due to small inorganic ions from dissociated strong electrolytes.¹ In a paper by Niels Bjerrum at the 16th Scandinavian Scientist Meeting in 1916, he concluded that the van't Hoff factor had erroneously been considered as a measure of the degree of dissociation of strong electrolytes.² Bjerrum proposed the strong electrolytes to be completely dissociated and he introduced an osmotic coefficient to account for the measured osmotic pressure.² The molal osmotic coefficient, ϕ , was expressed by Robinson and Stokes in terms of water activity, defined as the vapour pressure of an aqueous solution relative to that of pure water.³ We proposed the osmotic coefficient to be a product of two factors, $\phi = \phi^d \phi^w$.⁴ The term ϕ^d depends on the degree of dissociation of the dissolved strong electrolyte (NaCl, NH₄Cl, Na₂SO₄ or (NH₄)₂SO₄) and ϕ^w depends on changes in hydrogen bonded water clusters with solute concentration.⁴ The ϕ^w -term was measured from intensity changes of a band at 180 cm⁻¹ in the $R(\bar{\nu})$ -representation of the Raman spectrum.⁴ This contribution presents results for aqueous solution of KCl and MgCl₂.




Figure 1: Osmotic coefficients and degree of dissociation for KCl.

Raman spectra were recorded as previously described.⁴ For KCl the area of the 180 cm⁻¹ band was used to calculate the amount of tetrahedral bound water. The ratio between the amount of "free" water in pure water relative to that in a solution defines ϕ^w . Subsequently ϕ^d was calculated from values of the osmotic coefficient, ϕ , given by Hamer and Wu.⁵ See results in Figure 1. The minimum in ϕ around 0.5 molal is caused by a combination of changes in water structure, ϕ^w , and electrolyte dissociation, ϕ^d . At high

concentration the degree of dissociation, α , approaches 0.65, thus KCl is not completely dissociated. Similar results were obtained for MgCl₂ solutions, where also the strong H₂O---Mg²⁺ binding was taken into account. We hope the results will lead to a better understanding of osmolarity/osmolality on a molecular scale.

References

- [1] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, P. Walter, *Molecular Biology of the Cell* 2008, 5th ed., Garland Science, New York, US and Abingdon, UK, pp. 663-664
- [2] N. Bjerrum, *Fysisk Tidsskrift* 1916-1917, **15**, 59-73: Translated to English in *Niels Bjerrum, Selected Papers*, eds. N. Bohr *et al.*, Munksgaard, Copenhagen, Denmark, 1949 pp. 58-70.
- [3] R.A. Robinson, R.H. Stokes, *Electrolyte Solutions* 1959, 2nd ed., Butterworth, London, UK, pp. 25 and 29.
- [4] M. Frosch, M. Bilde, O.F. Nielsen, *J. Phys. Chem. A* 2010, **114**, pp. 11933-11942.
- [5] W.J. Hamer, Y.-C. Wu, *J. Phys. Chem. Ref. Data* 1972, **1**, 1047-1099.